




REPLANTATION

Patient Information Brochure

What is it?

"Replantation" refers to the surgical reattachment of a finger, hand, or arm that has been completely cut from a person's body (see Figure 1). The goal of replantation surgery is to give the patient back as much use of the injured area as possible. In some cases, replantation is not possible because the part is too damaged. If the lost part cannot be reattached, a patient may have to use a prosthesis (a device that substitutes for a missing part of the body). In many cases, a prosthesis may give a person without hands or arms the ability to function better than they would without the prosthesis.



Replantation is usually recommended when the replanted part will work at least as well as a prosthesis. Generally, a missing hand or finger would not be replanted knowing that it would not work, be painful, or get in the way of everyday life. Before surgery the doctor, if possible, will explain the procedure and how much use is likely to return following replantation. The patient or family member must decide whether that amount of use justifies the long and difficult operation, time

in the hospital, and months or years of rehabilitation.

How is the procedure done?

There are a number of steps in the replantation process. First, damaged tissue is carefully removed. Then bone ends are shortened and re-joined with pins or plates. This holds the part in place to allow the rest of the tissues to be restored to a normal position. Muscles, tendons, arteries, nerves and veins are then repaired.

What kind of recovery can I expect?

The patient has the most important role in the recovery process. Smoking causes poor circulation and may cause loss of blood flow to the replanted part. Allowing the replanted part to hang below heart level may also cause poor circulation. Younger patients have a better chance of their nerves growing back; they may regain more feeling and may regain more movement in the replanted part.

Generally, the further down the arm the injury occurs, the better the return of use of the replanted part to the patient. Patients who have not injured a joint will get more movement back than those with a joint injury. A cleanly cut part usually works better after replantation than one that has been crushed or pulled off. Recovery of use depends on re-growth of two types of nerves: sensory nerves that let you feel, and motor nerves that tell your muscles to move. Nerves grow about an inch per month. The number of inches from the injury to the tip of a finger gives the minimum number of months after which the patient may be able to feel something with that fingertip. The replanted part never regains 100% of its original use,



and most doctors consider 60% to 80% of use an excellent result.

Cold weather may be uncomfortable and provide reason for frequent complaint even for those with excellent recovery.

What about therapy and rehabilitation?

Complete healing of the injury and surgical wounds is only the beginning of a long process of rehabilitation. Therapy and temporary bracing are important to the recovery process. From the beginning, braces are used to protect the newly repaired tendons but allow the patient to move the replanted part. Therapy with limited motion helps keep joints from getting stiff, helps keep muscles mobile, and helps keep scar tissue to a minimum. Even after you have recovered, you may find that you cannot do everything you wish to do. Tailor-made devices may help many patients do special activities or hobbies. Talk to your physician or therapist to find out more about such devices. Many replant patients are able to return to the jobs they held before the injury. When this is not possible, patients can seek assistance in selecting a new type of work.

Are emotional problems common following replantation?

Replantation can affect your emotional life as well as your body. When your bandages are removed, and you see the replanted part for the first time, you may feel shock, grief, anger, disbelief, or disappointment because the replanted part simply does not look like it did before. Worries about the look of a replanted part and how it will work are common. Talking about these feelings with your doctor often helps you come to terms

REPLANTATION

Patient Information Brochure

with the outcome of the replantation. Your doctor may also ask a counsellor to assist with this process. You may find it helpful to talk with someone about it, and work through your feelings so you can move on with your life.

Will additional surgery be necessary?

After replantation surgery, some patients may need additional surgery at a later time to gain better function of the part. Some of the more common procedures are:

- Tenolysis: Frees tendons from scar tissue.
- Capsulotomy: Releases stiff, locked joints.
- Tendon or muscle transfer: Moves tendons or muscles to another spot so that they can work in an area that needs the tendon or muscle more.
- Nerve grafting: Replaces a scarred nerve or a gap in the nerves to improve how the nerve works.
- Late amputation: Removing the part because it does not work well, interferes with use of the hand, or has become painful.

Stay in the flow of life. You have many great gifts. Even with the best medical care, you need to be strong during the course of recovery. Remember that quality of life is directly related to your attitude and expectations - not on just regaining limb use.

The above material has been modified from content available at the American Society for Surgery of the Hand (ASSH) website. It is copyrighted by ASSH and its use is permitted for members.